



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C07H 21/04, A61K 48/00, C07H 21/02, C12Q 1/68, XXXX XXX/XXXXX	A1	(11) International Publication Number: WO 00/40595 (43) International Publication Date: 13 July 2000 (13.07.2000)
(21) International Application Number: PCT/US99/29593 (22) International Filing Date: 14 December 1999 (14.12.1999) (30) Priority Data: 09/226,568 07 January 1999 (07.01.1999) US (60) Parent Application or Grant ISIS PHARMACEUTICALS, INC. [/]; (). ACKERMANN, Elizabeth, J. [/]; (). BENNETT, C., Frank [/]; (). DEAN, Nicholas, M. [/]; (). MARCUSSON, Eric, G. [/]; (). ACKERMANN, Elizabeth, J. [/]; (). BENNETT, C., Frank [/]; (). DEAN, Nicholas, M. [/]; (). MARCUSSON, Eric, G. [/]; (). LICATA, Jane, Massey ; ().		Published
(54) Title: ANTISENSE MODULATION OF NOVEL ANTI-APOPTOTIC BCL-2-RELATED PROTEINS (54) Titre: MODULATION ANTISENS DE PROTEINES ANTI-APOPTOSE PROCHES DE BCL-2EINS (57) Abstract <p>Compositions and methods are provided for modulating the expression of novel anti-apoptotic bcl-2-related proteins. Antisense oligonucleotides targeted to nucleic acids encoding the human novel anti-apoptotic bcl-2-related proteins A1 and mcl-1 are preferred. Methods of using these compounds for modulation of novel anti-apoptotic bcl-2-related protein expression and for treatment of diseases associated with expression of novel anti-apoptotic bcl-2-related proteins are also provided. Also provided are methods of using these compounds for promoting apoptosis and for treatment of diseases for which promotion of apoptosis is desired.</p> (57) Abrégé <p>L'invention concerne des compositions et des techniques qui permettent de moduler l'expression de nouvelles protéines anti-apoptose proches de bcl-2. Les composés préférés sont des oligonucléotides antisens A1 et mcl-1 qui ciblent des acides nucléiques codant pour des protéines anti-apoptose proches de bcl-2. L'invention concerne également des méthodes qui permettent d'utiliser ces composés pour moduler l'expression des protéines anti-apoptose proches de bcl-2 et pour traiter les maladies associées à l'expression desdites protéines; ainsi que des méthodes qui permettent d'utiliser ces composés pour induire l'apoptose et traiter les maladies dans lesquelles on désire induire l'apoptose.</p>		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07H 21/04, 21/02, C12Q 1/68, 15/63, A61K 48/00		A1	(11) International Publication Number: WO 00/40595
			(43) International Publication Date: 13 July 2000 (13.07.00)
(21) International Application Number: PCT/US99/29593		(74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).	
(22) International Filing Date: 14 December 1999 (14.12.99)			
(30) Priority Data: 09/226,568 7 January 1999 (07.01.99) US		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/226,568 (CIP) Filed on 7 January 1999 (07.01.99)		Published With international search report.	
(71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): ACKERMANN, Elizabeth, J. [US/US]; 519 Santa Victoria, Solana Beach, CA 92075 (US). BENNETT, C., Frank [US/US]; 1347 Cassins Street, Carlsbad, CA 92008 (US). DEAN, Nicholas, M. [GB/US]; 2110 Whisperwind Lane, Olivenhain, CA 92024 (US). MARCUSSON, Eric, G. [US/US]; 6369 Caminito de Pastel, San Diego, CA 92111 (US).			
(54) Title: ANTISENSE MODULATION OF NOVEL ANTI-APOPTOTIC BCL-2-RELATED PROTEINS			
(57) Abstract <p>Compositions and methods are provided for modulating the expression of novel anti-apoptotic bcl-2-related proteins. Antisense oligonucleotides targeted to nucleic acids encoding the human novel anti-apoptotic bcl-2-related proteins A1 and mcl-1 are preferred. Methods of using these compounds for modulation of novel anti-apoptotic bcl-2-related protein expression and for treatment of diseases associated with expression of novel anti-apoptotic bcl-2-related proteins are also provided. Also provided are methods of using these compounds for promoting apoptosis and for treatment of diseases for which promotion of apoptosis is desired.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Description

5

10

15

20

25

30

35

40

45

50

55

THIS PAGE BLANK (USPTO)

- 1 -

ANTISENSE MODULATION OF NOVEL ANTI-APOPTOTIC
BCL-2-RELATED PROTEINS

FIELD OF THE INVENTION

The present invention provides compositions and methods for modulating the expression of novel anti-apoptotic bcl-2-related proteins. In particular, this invention relates to antisense compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding anti-apoptotic human bcl-2-related proteins. Such oligonucleotides have been shown to modulate the expression of novel anti-apoptotic bcl-2-related proteins.

BACKGROUND OF THE INVENTION

Programmed cell death, or apoptosis, is an essential feature of growth and development, as the control of cell number is a balance between cell proliferation and cell death. Apoptosis is an active rather than a passive process, resulting in cell suicide as a result of any of a number of external or internal signals. Apoptotic cell death is characterized by nuclear condensation, endonucleolytic degradation of DNA at nucleosomal intervals ("laddering") and plasma membrane blebbing. Programmed cell death plays an essential role in, for example, immune system development and nervous system development. In the former, T cells displaying autoreactive antigen receptors are removed by apoptosis. In the latter, a significant reshaping of neural structures occurs, partly through apoptosis.

An increasing number of genes and gene products have been implicated in apoptosis. One of these is bcl-2, which is an intracellular membrane protein shown to block or delay apoptosis. Overexpression of bcl-2 has been shown to be

- 2 -

5 related to hyperplasia, autoimmunity and resistance to
apoptosis, including that induced by chemotherapy (Fang et
al., *J. Immunol.* 1994, 153, 4388-4398). A family of bcl-2-
10 related genes has been described. All bcl-2 family members
5 share two highly conserved domains, BH1 and BH2. bcl-2 family
members include, but are not limited to, A1, mcl-1, bcl-w,
bax, bad, bak and bcl-x. A1, mcl-1, bcl-w and bcl-xl (long
15 form of bcl-x) are presently known to confer protection
against apoptosis and are referred to herein as "anti-
10 apoptotic bcl-2-related proteins." Of these, A1 and mcl-1 are
known as "novel anti-apoptotic bcl-2-related proteins." In
contrast, bax, bad, bak and bcl-xs (short form of bcl-x) are
20 presently known to promote cell death by inhibiting this
protective effect. The present invention relates to the novel
15 anti-apoptotic human bcl-2-related proteins, particularly
human A1 and mcl-1, and inhibition of the expression of these
proteins using antisense technology.

The gene encoding A1 (also known as bcl-2-related gene
30 expressed in fetal liver, or bfl-1, and Glasgow Rearranged
20 Sequence, or GRS) was identified as an early response gene in
murine hematopoietic cells treated with granulocyte-macrophage
colony-stimulating factor. The human homolog was subsequently
35 found to have extensive homology to bcl-2, especially within
the BH1 and BH2 domains. A correlation was noted between the
25 expression level of A1 and the development of stomach cancer
in clinical samples. Choi et al., 1995, *Oncogene* 11, 1693-
40 1698; WO 96/30513. The coding sequence of human A1 was cloned
(Karsan et al., 1996, *Blood* 87, 3089-3096; Genbank accession
no. U29680) and found to be expressed by hematopoietic cells
45 30 but also a variety of nonhematopoietic tissues. A1 is rapidly
inducible by phorbol esters and inflammatory cytokines, and
possibly by vascular endothelial growth factor. A1 is
believed to play a role in the regulation of physiological
50 cell death during embryonic development. Carrio et al., 1996,
35 *Am. J. Path.*, 149, 2133-2142. There is slight variation among

- 3 -

5 A1 sequences, with the human A1 sequence originally defined
as "GRS" differing by two amino acids from the A1 sequence
originally defined as "bfl-1". The GRS sequence was originally
10 isolated by NIH3T3 focus formation assay using DNA obtained
5 from a patient with chronic myeloid leukemia. Kenny et al,
Oncogene 14, 1997 997-1001. These researchers found a high
level of A1 expression in the cancer cell lines U-937
15 (histiocytic lymphoma), HL-60 (promyelocytic leukemia) and
Raji (Burkitt lymphoma) cells. Expression of A1 was also
20 found in THP-1 (acute myeloid leukemia), BJAB (Burkitt
lymphoma), activated Jurkat (acute T-cell leukemia) cells and
K-562 erythroleukemia (CML blast crisis) cells.

mcl-1 was originally identified from the differentiating
human myeloid leukemia cell line ML-1. Its expression was
25 found to increase early in the induction or "programming" of
differentiation of ML-1 cells before the appearance of
differentiation markers. The coding region of mcl-1 was
sequenced and found to have a pronounced region of sequence
30 homology to bcl-2 in the carboxyl-terminal region. Kozopas et
al., *Proc. Natl. Acad. Sci. USA.*, 1993, 90, 3516-3520; Genbank
accession no. L08246. Unlike bcl-2, mcl-1 contains a strong
PEST sequence (enriched in proline, glutamic acid, serine and
35 threonine) which is present in a variety of proteins that
undergo rapid turnover.

25 Overexpression of exogenously introduced mcl-1 has been
shown to cause a prolongation of viability under conditions
40 that normally cause apoptotic cell death, such as exposure to
cytotoxic agents (the chemotherapeutic agent etoposide,
calcium ionophore or UV irradiation) or the withdrawal of
30 required growth factors. Zhou et al., *Blood* 89, 1997, 630-
643.

Diseases and conditions in which apoptosis has been
implicated fall into two categories, those in which there is
50 increased cell survival (i.e., apoptosis is reduced) and those
35 in which there is excess cell death (i.e., apoptosis is

55

- 4 -

5 increased). Diseases in which there is an excessive
accumulation of cells due to increased cell survival include
cancer, autoimmune disorders and viral infections. Until
10 recently, it was thought that cytotoxic drugs killed target
5 cells directly by interfering with some life-maintaining
function. However, of late, it has been shown that exposure
to several cytotoxic drugs with disparate mechanisms of action
15 induces apoptosis in both malignant and normal cells.
Manipulation of levels of trophic factors (e.g., by anti-
10 estrogen compounds or those which reduce levels of various
growth hormones) has been one clinical approach to promote
20 apoptosis, since deprivation of trophic factors can induce
apoptosis. Apoptosis is also essential for the removal of
potentially autoreactive lymphocytes during development and
25 the removal of excess cells after the completion of an immune
or inflammatory response. Recent work has clearly
demonstrated that improper apoptosis may underlie the
pathogenesis of autoimmune diseases by allowing abnormal
30 autoreactive lymphocytes to survive. For these and other
20 conditions in which insufficient apoptosis is believed to be
involved, promotion of apoptosis is desired. Inhibition of
novel anti-apoptotic bcl-2-related proteins according to the
35 present invention is believed to result in promotion of
apoptosis.

25 In the second category, AIDS and neurodegenerative
disorders like Alzheimer's or Parkinson's disease represent
40 disorders for which an excess of cell death due to promotion
of apoptosis (or unwanted apoptosis) has been implicated.
Amyotrophic lateral sclerosis, retinitis pigmentosa, and
45 epilepsy are other neurologic disorders in which apoptosis has
been implicated. Apoptosis has been reported to occur in
conditions characterized by ischemia, e.g. myocardial
infarction and stroke. Apoptosis has also been implicated in
50 a number of liver disorders including obstructive jaundice and
35 hepatic damage due to toxins and drugs. Apoptosis has also

- 5 -

5 been identified as a key phenomenon in some diseases of the
kidney, i.e. polycystic kidney, as well as in disorders of the
pancreas including diabetes (Thatte, et al., *Drugs*, 1997, 54,
10 511-532). For these and other diseases and conditions in
5 which unwanted apoptosis is believed to be involved,
inhibitors of apoptosis are desired.

15 Antisense oligonucleotides have been used to elucidate
the role of several members of the bcl-2 family. Extensive
studies using antisense oligonucleotides targeted to bcl-2
10 have been performed, and an antisense compound (G3139, Genta
Incorporated) targeted to human bcl-2 has entered clinical
20 trials for lymphoma and prostate cancer.

Amarante-Mendes et al., *Oncogene*, 1998, 16, 1383-1390,
disclose antisense oligonucleotides targeted to bcr and bcl-x.
25 15 The latter downregulated the expression of bcl-xl and
increased the susceptibility of HL-60 Bcr-Abl cells to
staurosporine.

U.S. Patent 5,583,034 (Green et al.) discloses antisense
30 oligonucleotides which hybridize to the nucleic acid sequence
20 of an anti-apoptotic gene, preferably to the translation start
site of bcr-abl.

Wang et al. used a phosphorothioate oligonucleotide
35 targeted to the bcl-x translation start site to block CD40L-
mediated apoptotic rescue in murine WEHI-231 lymphoma cells
25 (*J. Immunol.*, 1995, 155, 3722-3725).

40 Fujio et al. have used an antisense oligodeoxynucleotide
targeted to murine and rat bcl-x mRNA to reduce bcl-xl protein
expression (*J. Clin. Invest.*, 1997, 99, 2898-2905). The
compound tested was the same as that of Wang et al.
45 30 Oligonucleotide treatment inhibited the cytoprotective effect
of leukemia inhibitory factor in mouse or rat cardiac
myocytes.

50 Pollman et al. used antisense oligodeoxynucleotides with
phosphorothioate backbones to downregulate bcl-xl expression
35 in blood vessel intimal cells (*Nature Med.*, 1998, 4, 222-227).

55

- 6 -

5 This resulted in induction of apoptosis and regression of
vascular lesions. Antisense sequences were targeted to the
translation initiation codon of mouse/human bcl-x (conserved
10 sequence) and were used in rabbits. Gibbons et al., U.S.
5 Patent 5,776,905, disclose methods for targeted deletion of
intimal lesion cells in the vasculature of a mammal with
vascular disease, preferably with antisense molecules specific
15 for anti-apoptotic genes, more preferably bcl-x and most
preferably bcl-xl.

10 Thompson et al., U.S. Patent 5,646,008 and WO 95/00642
describe an isolated and purified polynucleotide that encodes
a polypeptide other than bcl-2 that promotes or inhibits
programmed vertebrate cell death. Preferably the polypeptide
is bcl-xl, bcl-xs or bcl-x_l. Polypeptides, polynucleotides
25 identical or complementary to a portion of the isolated and
purified polynucleotide, expression vectors, host cells,
antibodies and therapeutic and diagnostic methods of use are
also provided.

30 Yang et al., WO 98/05777 disclose bcl-xy (gamma), a
20 novel isoform of the bcl-x family which includes an ankyrin
domain. Polypeptide and nucleic acid sequences for this
isoform are disclosed, as well as, *inter alia*, methods for
35 modulating bcl-xy activity, including antisense methods.

Chao et al., *Molec. Cell. Biol.*, 1998, 18, 4883-4898,
25 used an antisense construct containing the entire human mcl-1
cDNA in antisense orientation to show that down-regulation of
40 endogenous mcl-1 in TF-1 cells can induce apoptosis of these
cells.

Cory et al. (WO 97/35971) disclose methods for
30 modulating expression of bcl-w in a mammal by contacting the
bcl-w gene with an effective amount of a modulator of bcl-w
expression. Both enhanced and decreased bcl-w expression,
including use of antisense sequences to bcl-w, are disclosed.

50

55

- 7 -

WO 96/30513 (Shin et al.) discloses the sequence of a Bcl-2 related gene, Bfl-1, and use of a Bcl-2 related gene for diagnosing cancer.

WO 94/29330 discloses, inter alia, the mcl-1 polypeptide sequence and polynucleotide sequence, host cells and vectors containing the latter, antibodies which bind to the mcl-1 polypeptide, and diagnostic and therapeutic methods using these compounds. Methods for treating a subject with a mcl-1 associated cell proliferative disorder are generally disclosed, including antisense oligonucleotide and ribozyme approaches. No specific sequence or targeting information is provided.

SUMMARY OF THE INVENTION

The present invention is directed to antisense compounds, particularly oligonucleotides, which are targeted to a nucleic acid encoding novel anti-apoptotic bcl-2-related proteins, and which modulate the expression of these family members. In preferred embodiments the antisense oligonucleotides are targeted to human A1 or human mcl-1. Pharmaceutical compositions comprising the antisense compounds of the invention are also provided. Further provided are methods of modulating the expression of novel anti-apoptotic bcl-2-related proteins in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Methods for promoting apoptosis in cells or tissues are also provided. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of novel anti-apoptotic bcl-2-related proteins, or for which an increase in apoptosis is desired, by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

- 8 -

DETAILED DESCRIPTION OF THE INVENTION

The present invention comprehends antisense compounds capable of modulating expression of novel anti-apoptotic bcl-2-related proteins, particularly human A1 and mcl-1. A1 and mcl-1 inhibit apoptosis and therefore inhibitors of these targets are desired as promoters of apoptosis.

The present invention employs oligomeric antisense compounds, particularly oligonucleotides, for use in modulating the function of nucleic acid molecules encoding novel anti-apoptotic bcl-2-related proteins, ultimately modulating the amount of bcl-2-related protein produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding novel anti-apoptotic bcl-2-related proteins. As used herein, the terms "target nucleic acid" and "nucleic acid encoding a novel anti-apoptotic bcl-2-related protein" encompass DNA encoding a novel anti-apoptotic bcl-2-related protein, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of the novel anti-apoptotic bcl-2-related protein. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of

5 a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

10 It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding a novel anti-apoptotic bcl-2-related protein. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for

- 10 -

translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used *in vivo* to initiate translation of an mRNA molecule transcribed from a gene encoding a novel anti-apoptotic bcl-2-related protein, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, *i.e.*, 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (*i.e.*, 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (*i.e.*, 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding

- 11 -

5 nucleotides on the gene. The 5' cap of an mRNA comprises an
N7-methylated guanosine residue joined to the 5'-most residue
of the mRNA via a 5'-5' triphosphate linkage. The 5' cap
10 region of an mRNA is considered to include the 5' cap
5 structure itself as well as the first 50 nucleotides adjacent
to the cap. The 5' cap region may also be a preferred target
region.

15 Although some eukaryotic mRNA transcripts are directly
translated, many contain one or more regions, known as
10 "introns," which are excised from a transcript before it is
translated. The remaining (and therefore translated) regions
20 are known as "exons" and are spliced together to form a
continuous mRNA sequence. mRNA splice sites, i.e., intron-
exon junctions, may also be preferred target regions, and are
25 particularly useful in situations where aberrant splicing is
implicated in disease, or where an overproduction of a
particular mRNA splice product is implicated in disease.
Aberrant fusion junctions due to rearrangements or deletions
30 are also preferred targets. It has also been found that
20 introns can also be effective, and therefore preferred, target
regions for antisense compounds targeted, for example, to DNA
or pre-mRNA.

35 Once one or more target sites have been identified,
oligonucleotides are chosen which are sufficiently
25 complementary to the target, i.e., hybridize sufficiently well
and with sufficient specificity, to give the desired effect.

40 In the context of this invention, "hybridization" means
hydrogen bonding, which may be Watson-Crick, Hoogsteen or
reversed Hoogsteen hydrogen bonding, between complementary
45 30 nucleoside or nucleotide bases. For example, adenine and
thymine are complementary nucleobases which pair through the
formation of hydrogen bonds. "Complementary," as used herein,
refers to the capacity for precise pairing between two
50 nucleotides. For example, if a nucleotide at a certain
35 position of an oligonucleotide is capable of hydrogen bonding

- 12 -

5 with a nucleotide at the same position of a DNA or RNA
molecule, then the oligonucleotide and the DNA or RNA are
considered to be complementary to each other at that position.
10 The oligonucleotide and the DNA or RNA are complementary to
5 each other when a sufficient number of corresponding positions
in each molecule are occupied by nucleotides which can
hydrogen bond with each other. Thus, "specifically
15 hybridizable" and "complementary" are terms which are used to
indicate a sufficient degree of complementarity or precise
10 pairing such that stable and specific binding occurs between
the oligonucleotide and the DNA or RNA target. It is
20 understood in the art that the sequence of an antisense
compound need not be 100% complementary to that of its target
nucleic acid to be specifically hybridizable. An antisense
25 compound is specifically hybridizable when binding of the
compound to the target DNA or RNA molecule interferes with the
normal function of the target DNA or RNA to cause a loss of
utility, and there is a sufficient degree of complementarity
30 to avoid non-specific binding of the antisense compound to
20 non-target sequences under conditions in which specific
binding is desired, i.e., under physiological conditions in
the case of *in vivo* assays or therapeutic treatment, and in
35 the case of *in vitro* assays, under conditions in which the
assays are performed.

25 Antisense compounds are commonly used as research
reagents and diagnostics. For example, antisense
40 oligonucleotides, which are able to inhibit gene expression
with exquisite specificity, are often used by those of
ordinary skill to elucidate the function of particular genes.
45 30 Antisense compounds are also used, for example, to distinguish
between functions of various members of a biological pathway.
Antisense modulation has, therefore, been harnessed for
research use.

50 The specificity and sensitivity of antisense is also
35 harnessed by those of skill in the art for therapeutic uses.

55

- 13 -

5 Antisense oligonucleotides have been employed as therapeutic
moieties in the treatment of disease states in animals and
man. Antisense oligonucleotides have been safely and
10 effectively administered to humans and numerous clinical
5 trials are presently underway. It is thus established that
oligonucleotides can be useful therapeutic modalities that can
be configured to be useful in treatment regimes of cells,
15 tissues and animals, especially humans. In the context of
this invention, the term "oligonucleotide" refers to an
10 oligomer or polymer of ribonucleic acid (RNA) or
deoxyribonucleic acid (DNA) or mimetics thereof. This term
20 includes oligonucleotides composed of naturally-occurring
nucleobases, sugars and covalent internucleoside (backbone)
linkages as well as oligonucleotides having non-naturally-
25 occurring portions which function similarly. Such modified
or substituted oligonucleotides are often preferred over
native forms because of desirable properties such as, for
example, enhanced cellular uptake, enhanced affinity for
30 nucleic acid target and increased stability in the presence
20 of nucleases.

While antisense oligonucleotides are a preferred form
of antisense compound, the present invention comprehends other
35 oligomeric antisense compounds, including but not limited to
oligonucleotide mimetics such as are described below. The
25 antisense compounds in accordance with this invention
preferably comprise from about 8 to about 30 nucleobases.
40 Particularly preferred are antisense oligonucleotides
comprising from about 8 to about 30 nucleobases (*i.e.* from
about 8 to about 30 linked nucleosides). As is known in the
30 art, a nucleoside is a base-sugar combination. The base
45 portion of the nucleoside is normally a heterocyclic base.
The two most common classes of such heterocyclic bases are the
purines and the pyrimidines. Nucleotides are nucleosides that
50 further include a phosphate group covalently linked to the
35 sugar portion of the nucleoside. For those nucleosides that

- 14 -

5 include a pentofuranosyl sugar, the phosphate group can be
linked to either the 2'-, 3'- or 5'- hydroxyl moiety of the
sugar. In forming oligonucleotides, the phosphate groups
10 covalently link adjacent nucleosides to one another to form
5 a linear polymeric compound. In turn, the respective ends of
this linear polymeric structure can be further joined to form
a circular structure. However, open linear structures are
15 generally preferred. Within the oligonucleotide structure,
the phosphate groups are commonly referred to as forming the
10 internucleoside backbone of the oligonucleotide. The normal
linkage or backbone of RNA and DNA is a 3' to 5'
20 phosphodiester linkage.

Specific examples of preferred antisense compounds
useful in this invention include oligonucleotides containing
25 15 modified backbones or non-natural internucleoside linkages.
As defined in this specification, oligonucleotides having
modified backbones include those that retain a phosphorus atom
in the backbone and those that do not have a phosphorus atom
30 in the backbone. For the purposes of this specification, and
20 as sometimes referenced in the art, modified oligonucleotides
that do not have a phosphorus atom in their internucleoside
backbone can also be considered to be oligonucleosides.

35 Preferred modified oligonucleotide backbones include,
for example, phosphorothioates, chiral phosphorothioates,
25 phosphorodithioates, phosphotriesters, aminoalkylphosphotri-
esters, methyl and other alkyl phosphonates including 3'-
40 alkylene phosphonates and chiral phosphonates, phosphinates,
phosphoramidates including 3'-amino phosphoramidate and
aminoalkylphosphoramidates, thionophosphoramidates, thiono-
45 30 alkylphosphonates, thionoalkylphosphotriesters, and borano-
phosphates having normal 3'-5' linkages, 2'-5' linked analogs
of these, and those having inverted polarity wherein the
adjacent pairs of nucleoside units are linked 3'-5' to 5'-3'
50 or 2'-5' to 5'-2'. Various salts, mixed salts and free acid
35 forms are also included.

- 15 -

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of

- 16 -

the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular $-\text{CH}_2-\text{NH}-\text{O}-\text{CH}_2-$, $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{O}-\text{CH}_2-$ [known as a methylene (methylimino) or MMI backbone], $-\text{CH}_2-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-$, $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{N}(\text{CH}_3)-\text{CH}_2-$ and $-\text{O}-\text{N}(\text{CH}_2)_f\text{CH}-\text{CH}_2-$ [wherein the native phosphodiester backbone is represented as $-\text{O}-\text{P}-\text{O}-\text{CH}_2-$] of the above referenced U.S. Patent 5,489,677, and the amide backbones of the above referenced U.S. Patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. Patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C_1 to C_{10} alkyl or C_2 to C_{10} alkenyl and alkynyl. Particularly preferred are $\text{O}[(\text{CH}_2)_n\text{O}]_m\text{CH}_3$, $\text{O}(\text{CH}_2)_n\text{OCH}_3$, $\text{O}(\text{CH}_2)_n\text{NH}_2$, $\text{O}(\text{CH}_2)_n\text{CH}_3$, $\text{O}(\text{CH}_2)_n\text{ONH}_2$,

- 17 -

5 and $O(CH_2)_nON[(CH_2)_mCH_3]$, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C_1 to C_{10} lower alkyl, 10 substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH_3 , OCN, Cl, Br, CN, CF_3 , OCF_3 , $SOCH_3$, SO_2CH_3 , ONO_2 , NO_2 , N_3 , NH_2 , heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA 15 cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes an alkoxyalkoxy group, 2'-methoxyethoxy ($2'-O-CH_2CH_2OCH_3$, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., 25 *Helv. Chim. Acta*, 1995, 78, 486-504). A further preferred modification includes 2'-dimethylaminooxyethoxy, i.e., a $O(CH_2)_nON(CH_3)_2$ group, also known as 2'-DMAOE.

Other preferred modifications include 2'-methoxy ($2'-O-CH_3$), 2'-aminopropoxy ($2'-OCH_2CH_2CH_2NH_2$) and 2'-fluoro ($2'-F$). 20 Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics 25 such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 30 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, each of which is herein incorporated by reference.

50 Oligonucleotides may also include nucleobase (often 35 referred to in the art simply as "base") modifications or

- 18 -

substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Patent 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pp. 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie, International Edition*, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Crooke, S.T., and Lebleu, B. eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 289-302. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyl-adenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more

- 19 -

particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; 5,681,941; and 5,750,692, each of which is herein incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36,

- 20 -

3651-3654), a palmitoyl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937).

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional

- 21 -

5 region of the oligonucleotide may serve as a substrate for
enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By
10 way of example, RNase H is a cellular endonuclease which
cleaves the RNA strand of an RNA:DNA duplex. Activation of
5 RNase H, therefore, results in cleavage of the RNA target,
thereby greatly enhancing the efficiency of oligonucleotide
inhibition of gene expression. Cleavage of the RNA target can
15 be routinely detected by gel electrophoresis and, if
necessary, associated nucleic acid hybridization techniques
10 known in the art.

20 Chimeric antisense compounds of the invention may be
formed as composite structures of two or more
oligonucleotides, modified oligonucleotides, oligonucleosides
and/or oligonucleotide mimetics as described above. Such
25 compounds have also been referred to in the art as hybrids or
gapmers. Representative United States patents that teach the
preparation of such hybrid structures include, but are not
limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775;
30 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065;
20 5,652,355; 5,652,356; and 5,700,922, each of which is herein
incorporated by reference.

35 The antisense compounds used in accordance with this
invention may be conveniently and routinely made through the
well-known technique of solid phase synthesis. Equipment for
25 such synthesis is sold by several vendors including, for
example, Applied Biosystems (Foster City, CA). Any other
40 means for such synthesis known in the art may additionally or
alternatively be employed. It is well known to use similar
techniques to prepare oligonucleotides such as the
30 phosphorothioates and alkylated derivatives.

45 The antisense compounds of the invention are synthesized
in vitro and do not include antisense compositions of
biological origin, or genetic vector constructs designed to
50 direct the *in vivo* synthesis of antisense molecules.

- 22 -

5 The compounds of the invention may also be admixed,
encapsulated, conjugated or otherwise associated with other
molecules, molecule structures or mixtures of compounds, as
10 for example, liposomes, receptor targeted molecules, oral,
5 rectal, topical or other formulations, for assisting in
uptake, distribution and/or absorption. Representative United
States patents that teach the preparation of such uptake,
15 distribution and/or absorption assisting formulations include,
but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016;
10 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020;
5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921;
20 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619;
5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295;
5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and
25 15 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any
pharmaceutically acceptable salts, esters, or salts of such
esters, or any other compound which, upon administration to
30 an animal including a human, is capable of providing (directly
20 or indirectly) the biologically active metabolite or residue
thereof. Accordingly, for example, the disclosure is also
drawn to prodrugs and pharmaceutically acceptable salts of the
35 compounds of the invention, pharmaceutically acceptable salts
of such prodrugs, and other bioequivalents.

25 The term "prodrug" indicates a therapeutic agent that
is prepared in an inactive form that is converted to an active
40 form (i.e., drug) within the body or cells thereof by the
action of endogenous enzymes or other chemicals and/or
conditions. In particular, prodrug versions of the
45 30 oligonucleotides of the invention are prepared as SATE
[(S-acetyl-2-thioethyl) phosphate] derivatives according to
the methods disclosed in WO 93/24510 or in WO 94/26764.

The term "pharmaceutically acceptable salts" refers to
50 physiologically and pharmaceutically acceptable salts of the
35 compounds of the invention: i.e., salts that retain the

- 23 -

desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred addition salts are acid salts such as the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid,

- 24 -

5 hydroxymaleic acid, methylmaleic acid, fumaric acid, malic
acid, tartaric acid, lactic acid, oxalic acid, gluconic acid,
10 glucaric acid, glucuronic acid, citric acid, benzoic acid,
cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic
5 acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embolic
acid, nicotinic acid or isonicotinic acid; and with amino
acids, such as the 20 alpha-amino acids involved in the
15 synthesis of proteins in nature, for example glutamic acid or
aspartic acid, and also with phenylacetic acid,
10 methanesulfonic acid, ethanesulfonic acid,
2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid,
20 benzenesulfonic acid, 4-methylbenzenesulfoic acid,
naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid,
2- or 3-phosphoglycerate, glucose-6-phosphate,
25 N-cyclohexylsulfamic acid (with the formation of cyclamates),
or with other acid organic compounds, such as ascorbic acid.
Pharmaceutically acceptable salts of compounds may also be
prepared with a pharmaceutically acceptable cation. Suitable
30 pharmaceutically acceptable cations are well known to those
skilled in the art and include alkaline, alkaline earth,
ammonium and quaternary ammonium cations. Carbonates or
hydrogen carbonates are also possible.

35 For oligonucleotides, preferred examples of
pharmaceutically acceptable salts include but are not limited
25 to (a) salts formed with cations such as sodium, potassium,
ammonium, magnesium, calcium, polyamines such as spermine and
40 spermidine, etc.; (b) acid addition salts formed with
inorganic acids, for example hydrochloric acid, hydrobromic
acid, sulfuric acid, phosphoric acid, nitric acid and the
30 like; (c) salts formed with organic acids such as, for
45 example, acetic acid, oxalic acid, tartaric acid, succinic
acid, maleic acid, fumaric acid, gluconic acid, citric acid,
malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic
50 acid, alginic acid, polyglutamic acid, naphthalenesulfonic
35 acid, methanesulfonic acid, p-toluenesulfonic acid,

- 25 -

5 naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

10 The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of one or more novel anti-apoptotic bcl-2-related proteins is treated 10 by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds 20 and methods of the invention may also be useful prophylactically, e.g., to prevent or delay inflammation or tumor formation, for example.

30 The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding novel anti-apoptotic bcl-2-related proteins, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding a novel anti-apoptotic bcl-2-related protein can 35 be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of novel anti-apoptotic bcl-2-related 40 proteins in a sample may also be prepared.

45 The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of 50 ways depending upon whether local or systemic treatment is

55

- 26 -

5 desired and upon the area to be treated. Administration may
be topical (including ophthalmic and to mucous membranes
including vaginal and rectal delivery), pulmonary, e.g., by
10 inhalation or insufflation of powders or aerosols, including
5 by nebulizer; intratracheal, intranasal, epidermal and
transdermal), oral or parenteral. Parenteral administration
includes intravenous, intraarterial, subcutaneous,
15 intraperitoneal or intramuscular injection or infusion; or
intracranial, e.g., intrathecal or intraventricular,
10 administration. Oligonucleotides with at least one 2'-O-
methoxyethyl modification are believed to be particularly
20 useful for oral administration.

Pharmaceutical compositions and formulations for topical
administration may include transdermal patches, ointments,
25 lotions, creams, gels, drops, suppositories, sprays, liquids
and powders. Conventional pharmaceutical carriers, aqueous,
powder or oily bases, thickeners and the like may be necessary
or desirable. Coated condoms, gloves and the like may also
30 be useful.

20 Compositions and formulations for oral administration
include powders or granules, suspensions or solutions in water
or non-aqueous media, capsules, sachets or tablets.
35 Thickeners, flavoring agents, diluents, emulsifiers,
dispersing aids or binders may be desirable.

25 Compositions and formulations for parenteral,
intrathecal or intraventricular administration may include
40 sterile aqueous solutions which may also contain buffers,
diluents and other suitable additives such as, but not limited
to, penetration enhancers, carrier compounds and other
30 pharmaceutically acceptable carriers or excipients.

45 Pharmaceutical compositions and/or formulations
comprising the oligonucleotides of the present invention may
also include penetration enhancers in order to enhance the
50 alimentary delivery of the oligonucleotides. Penetration
35 enhancers may be classified as belonging to one of five broad

- 27 -

categories, i.e., fatty acids, bile salts, chelating agents, surfactants and non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, 8, 91-192; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). One or more penetration enhancers from one or more of these broad categories may be included.

Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, recinleate, monoolein (a.k.a. 1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arichidonic acid, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, mono- and di-glycerides and physiologically acceptable salts thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, 8:2, 91-192; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7:1, 1-33; El-Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654). Examples of some presently preferred fatty acids are sodium caprate and sodium laurate, used singly or in combination at concentrations of 0.5 to 5%.

The physiological roles of bile include the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 In: *Goodman & Gilman's The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al., eds., McGraw-Hill, New York, NY, 1996, pages 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus, the term "bile salt" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. A presently preferred bile salt is chenodeoxycholic acid (CDCA) (Sigma Chemical Company, St. Louis, MO), generally used at concentrations of 0.5 to 2%.

- 28 -

5 Complex formulations comprising one or more penetration
enhancers may be used. For example, bile salts may be used
in combination with fatty acids to make complex formulations.
10 Preferred combinations include CDCA combined with sodium
5 caprate or sodium laurate (generally 0.5 to 5%).

 Chelating agents include, but are not limited to,
disodium ethylenediaminetetraacetate (EDTA), citric acid,
15 salicylates (e.g., sodium salicylate, 5-methoxysalicylate and
homovanilate), N-acyl derivatives of collagen, laureth-9 and
10 N-amino acyl derivatives of beta-diketones (enamines) (Lee et
al., *Critical Reviews in Therapeutic Drug Carrier Systems*,
20 1991, 8:2, 92-192; Muranishi, *Critical Reviews in Therapeutic
Drug Carrier Systems*, 1990, 7:1, 1-33; Buur et al., *J. Control
Rel.*, 1990, 14, 43-51). Chelating agents have the added
25 15 advantage of also serving as DNase inhibitors.

 Surfactants include, for example, sodium lauryl sulfate,
polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl
ether (Lee et al., *Critical Reviews in Therapeutic Drug
Carrier Systems*, 1991, 8:2, 92-191); and perfluorochemical
30 20 emulsions, such as FC-43 (Takahashi et al., *J. Pharm.
Pharmacol.*, 1988, 40, 252-257).

 Non-surfactants include, for example, unsaturated cyclic
ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives
35 (Lee et al., *Critical Reviews in Therapeutic Drug Carrier
Systems*, 1991, 8:2, 92-191); and non-steroidal anti-
inflammatory agents such as diclofenac sodium, indomethacin
40 and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*,
1987, 39, 621-626).

 As used herein, "carrier compound" refers to a nucleic
45 30 acid, or analog thereof, which is inert (i.e., does not
possess biological activity per se) but is recognized as a
nucleic acid by in vivo processes that reduce the
bioavailability of a nucleic acid having biological activity
50 by, for example, degrading the biologically active nucleic

- 29 -

acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioated oligonucleotide in hepatic tissue is reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

In contrast to a carrier compound, a "pharmaceutically acceptable carrier" (excipient) is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The pharmaceutically acceptable carrier may be liquid or solid and is selected with the planned manner of administration in mind so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutically acceptable carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrates (e.g., starch, sodium starch glycolate, etc.); or wetting agents (e.g.,

- 30 -

5 sodium lauryl sulphate, etc.). Sustained release oral
delivery systems and/or enteric coatings for orally
administered dosage forms are described in U.S. Patents
10 4,704,295; 4,556,552; 4,309,406; and 4,309,404, which are
5 incorporated by reference.

The compositions of the present invention may
15 additionally contain other adjunct components conventionally
found in pharmaceutical compositions, at their art-established
usage levels. Thus, for example, the compositions may contain
10 additional compatible pharmaceutically-active materials such
as, e.g., antipruritics, astringents, local anesthetics or
20 anti-inflammatory agents, or may contain additional materials
useful in physically formulating various dosage forms of the
composition of present invention, such as dyes, flavoring
25 agents, preservatives, antioxidants, opacifiers, thickening
agents and stabilizers. However, such materials, when added,
should not unduly interfere with the biological activities of
the components of the compositions of the invention.

30 Regardless of the method by which the antisense
20 compounds of the invention are introduced into a patient,
colloidal dispersion systems may be used as delivery vehicles
to enhance the *in vivo* stability of the compounds and/or to
35 target the compounds to a particular organ, tissue or cell
type. Colloidal dispersion systems include, but are not
25 limited to, macromolecule complexes, nanocapsules,
microspheres, beads and lipid-based systems including oil-in-
40 water emulsions, micelles, mixed micelles, liposomes and
lipid:oligonucleotide complexes of uncharacterized structure.
A preferred colloidal dispersion system is a plurality of
45 liposomes. Liposomes are microscopic spheres having an
aqueous core surrounded by one or more outer layer(s) made up
of lipids arranged in a bilayer configuration (see, generally,
Chonn et al., *Current Op. Biotech.*, 1995, 6, 698-708).

50 Certain embodiments of the invention provide for
35 liposomes and other compositions containing (a) one or more

- 31 -

antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include, but are not limited to, anticancer drugs such as daunorubicin, dactinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine (CA), 5-fluorouracil (5-FU), floxuridine (5-FUdR), methotrexate (MTX), colchicine, vincristine, vinblastine, etoposide, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pp. 1206-1228. Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pp. 2499-2506 and 46-49, respectively. Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be

- 32 -

5 calculated from measurements of drug accumulation in the body
of the patient. Persons of ordinary skill can easily
determine optimum dosages, dosing methodologies and repetition
10 rates. Optimum dosages may vary depending on the relative
5 potency of individual oligonucleotides, and can generally be
estimated based on EC₅₀s found to be effective in in vitro and
in vivo animal models. In general, dosage is from 0.01 µg to
15 100 g per kg of body weight, and may be given once or more
daily, weekly, monthly or yearly, or even once every 2 to 20
10 years. Persons of ordinary skill in the art can easily
estimate repetition rates for dosing based on measured
20 residence times and concentrations of the drug in bodily
fluids or tissues. Following successful treatment, it may be
desirable to have the patient undergo maintenance therapy to
25 15 prevent the recurrence of the disease state, wherein the
oligonucleotide is administered in maintenance doses, ranging
from 0.01 µg to 100 g per kg of body weight, once or more
daily, to once every 20 years.

30 While the present invention has been described with
20 specificity in accordance with certain of its preferred
embodiments, the following examples serve only to illustrate
the invention and are not intended to limit the same.

35 EXAMPLES

Example 1

40 25 Nucleoside Phosphoramidites for Oligonucleotide Synthesis

Deoxy and 2'-alkoxy amidites

45 2'-Deoxy and 2'-methoxy beta-cyanoethyl-diisopropyl
phosphoramidites were purchased from commercial sources (e.g.
Chemgenes, Needham, MA or Glen Research, Inc. Sterling, VA).
30 Other 2'-O-alkoxy substituted nucleoside amidites are prepared
as described in U.S. Patent 5,506,351, herein incorporated by
50 reference. For oligonucleotides synthesized using 2'-alkoxy

- 33 -

amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods (Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling, VA or ChemGenes, Needham, MA).

2'-Fluoro amidites

2'-Fluorodeoxyadenosine amidites

2'-fluoro oligonucleotides were synthesized as described previously by Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841 and U.S. Patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S_N2 -displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

2'-Fluorodeoxyguanosine

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give

- 34 -

diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

2'-Fluorouridine

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

2'-Fluorodeoxycytidine

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

2'-O-(2-Methoxyethyl) modified amidites

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced

- 35 -

5 pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount
10 of methanol (ca. 400 mL). The solution was poured into fresh
5 ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 hours) to give a solid that was crushed to a light tan powder
15 (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt
10 (ca. 5%). The material was used as is for further reactions or purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C.

2'-O-Methoxyethyl-5-methyluridine

25 2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for
30 48 hours at 155-160°C, the vessel was opened and the solution
20 evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH₃CN (600 mL) and evaporated. A silica gel column (3 kg) was
35 packed in CH₂Cl₂/Acetone/MeOH (20:5:3) containing 0.5% Et₃NH.
25 The residue was dissolved in CH₂Cl₂ (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of
40 product. Additional material was obtained by reworking impure
45 30 fractions.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

50 2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue

55

- 36 -

5 dissolved in pyridine (1.3 L). A first aliquot of
dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the
mixture stirred at room temperature for one hour. A second
10 aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was
5 added and the reaction stirred for an additional one hour.
Methanol (170 mL) was then added to stop the reaction. HPLC
showed the presence of approximately 70% product. The solvent
15 was evaporated and triturated with CH₃CN (200 mL). The
residue was dissolved in CHCl₃ (1.5 L) and extracted with
10 2x500 mL of saturated NaHCO₃ and 2x500 mL of saturated NaCl.
The organic phase was dried over Na₂SO₄, filtered and
20 evaporated. 275 g of residue was obtained. The residue was
purified on a 3.5 kg silica gel column, packed and eluted with
EtOAc/Hexane/Acetone (5:5:1) containing 0.5% Et₃NH. The pure
25 fractions were evaporated to give 164 g of product.
Approximately 20 g additional was obtained from the impure
fractions to give a total yield of 183 g (57%).

30 3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-
methyluridine

20 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine
(106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture
35 prepared from 562 mL of DMF and 188 mL of pyridine) and acetic
anhydride (24.38 mL, 0.258 M) were combined and stirred at
room temperature for 24 hours. The reaction was monitored by
25 tlc by first quenching the tlc sample with the addition of
MeOH. Upon completion of the reaction, as judged by tlc, MeOH
(50 mL) was added and the mixture evaporated at 35°C. The
residue was dissolved in CHCl₃ (800 mL) and extracted with
45 2x200 mL of saturated sodium bicarbonate and 2x200 mL of
30 saturated NaCl. The water layers were back extracted with 200
mL of CHCl₃. The combined organics were dried with sodium
sulfate and evaporated to give 122 g of residue (approx. 90%
50 product). The residue was purified on a 3.5 kg silica gel
column and eluted using EtOAc/Hexane(4:1). Pure product

- 37 -

fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH_3CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH_3CN (1 L), cooled to -5°C and stirred for 0.5 hours using an overhead stirrer. POCl_3 was added dropwise, over a 30 minute period, to the stirred solution maintained at $0-10^\circ\text{C}$, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO_3 and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH_4OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH_3 gas was added and the vessel heated to 100°C for 2 hours (tlc showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The

- 38 -

organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, tlc showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl₃ (700 mL) and extracted with saturated NaHCO₃ (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO₄ and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/Hexane (1:1) containing 0.5% Et₃NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH₂Cl₂ (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (tlc showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO₃ (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH₂Cl₂ (300 mL), and the extracts were combined, dried over MgSO₄ and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/Hexane (3:1) as the eluting

- 39 -

solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

2'-O-(dimethylaminoxyethyl) nucleoside amidites

2'-(Dimethylaminoxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminoxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

5'-O-tert-Butyldiphenylsilyl-02-2'-anhydro-5-methyluridine

O²-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (Rf 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

- 40 -

5'-'O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine

10 In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In
5 the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-'O-tert-Butyldiphenylsilyl-O²-2'-
15 anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual
10 stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160°C was reached and then maintained for 16 h (pressure < 100 psig). The reaction
20 vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl
25 acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm
30 Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol.
20 [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The
35 residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate
25 fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield
40 based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure
30 product.

- 41 -

2'-O-([2-phthalimidoxy)ethyl]-5'-t-
butyldiphenylsilyl-5-methyluridine

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-
methyluridine (20g, 36.98mmol) was mixed with
5 triphenylphosphine (11.63g, 44.36mmol) and N-
hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over
15 P₂O₅ under high vacuum for two days at 40°C. The reaction
mixture was flushed with argon and dry THF (369.8mL, Aldrich,
sure seal bottle) was added to get a clear solution. Diethyl-
10 azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the
reaction mixture. The rate of addition is maintained such
that resulting deep red coloration is just discharged before
adding the next drop. After the addition was complete, the
reaction was stirred for 4 hrs. By that time TLC showed the
25 completion of the reaction (ethylacetate:hexane, 60:40). The
solvent was evaporated in vacuum. Residue obtained was placed
on a flash column and eluted with ethyl acetate:hexane
30 (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-
butyldiphenylsilyl-5-methyluridine as white foam (21.819,
20 86%).

5'-O-tert-butyldiphenylsilyl-2'-O-[(2-
formadoximinooxy)ethyl]-5-methyluridine_

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-
methyluridine (3.1g, 4.5mmol) was dissolved in dry CH₂Cl₂,
40 (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added
dropwise at -10°C to 0°C. After 1 hr the mixture was filtered,
the filtrate was washed with ice cold CH₂Cl₂, and the combined
organic phase was washed with water, brine and dried over
45 anhydrous Na₂SO₄. The solution was concentrated to get 2'-O-
30 (aminooxyethyl) thymidine, which was then dissolved in MeOH
(67.5mL). To this formaldehyde (20% aqueous solution, w/w,
1.1eg.) was added and the mixture for 1 hr. Solvent was
50 removed under vacuum; residue chromatographed to get 5'-O-

55

- 42 -

tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy) ethyl]-5-methyluridine as white foam (1.95, 78%).

5'-O-tert-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine

5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 hr, the reaction monitored by TLC (5% MeOH in CH₂Cl₂). Aqueous NaHCO₃ solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na₂SO₄, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO₃ (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na₂SO₄ and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH₂Cl₂ to get 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

- 43 -

2'-O-(dimethylaminoxyethyl)-5-methyluridine

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH₂Cl₂). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH₂Cl₂ to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P₂O₅ under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH₂Cl₂ (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P₂O₅ under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous

- 44 -

5 acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N',N'-
tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added.
10 The reaction mixture was stirred at ambient temperature for
4 hrs under inert atmosphere. The progress of the reaction
5 was monitored by TLC (hexane:ethyl acetate 1:1). The solvent
was evaporated, then the residue was dissolved in ethyl
15 acetate (70mL) and washed with 5% aqueous NaHCO₃ (40mL).
Ethyl acetate layer was dried over anhydrous Na₂SO₄ and
concentrated. Residue obtained was chromatographed (ethyl
20 acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-
dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-
diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

Example 2

Oligonucleotide Synthesis

25 Unsubstituted and substituted phosphodiester (P=O)
oligonucleotides are synthesized on an automated DNA
synthesizer (Applied Biosystems model 380B) using standard
30 phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as per the
20 phosphodiester oligonucleotides except the standard oxidation
bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-
35 3-one 1,1-dioxide in acetonitrile for the stepwise thiation
of the phosphite linkages. The thiation wait step was
increased to 68 seconds and was followed by the capping step.
25 After cleavage from the CPG column and deblocking in
concentrated ammonium hydroxide at 55°C (18 hr), the
oligonucleotides were purified by precipitating twice with 2.5
40 volumes of ethanol from a 0.5 M NaCl solution.

45 Phosphinate oligonucleotides are prepared as described
30 in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as
described in U.S. Patent 4,469,863, herein incorporated by
50 reference.

- 45 -

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

Example 3

Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

- 46 -

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

Example 4

PNA Synthesis

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

Example 5

Synthesis of Chimeric Oligonucleotides

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

- 47 -

[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric
Phosphorothioate Oligonucleotides

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 Ammonia/Ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hours at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hours at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate
Oligonucleotides

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

- 48 -

[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy
Phosphorothioate]--[2'-O-(2-Methoxyethyl)
Phosphodiester] Chimeric Oligonucleotides

[2'-O-(2-methoxyethyl phosphodiester)--[2'-deoxy phos-
phorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric
oligonucleotides are prepared as per the above procedure for
the 2'-O-methyl chimeric oligonucleotide with the substitution
of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites,
oxidization with iodine to generate the phosphodiester
internucleotide linkages within the wing portions of the
chimeric structures and sulfurization utilizing 3,4-dihydro-2H-
benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) to generate
the phosphorothioate internucleotide linkages for the center
gap.

Other chimeric oligonucleotides, chimeric oligonucleo-
sides and mixed chimeric oligonucleotides/oligonucleosides are
synthesized according to U.S. Patent 5,623,065, herein
incorporated by reference.

Example 6

Oligonucleotide Isolation

After cleavage from the controlled pore glass column
(Applied Biosystems) and deblocking in concentrated ammonium
hydroxide at 55°C for 18 hours, the oligonucleotides or
oligonucleosides were purified by precipitation twice out of
0.5 M NaCl with 2.5 volumes ethanol. Synthesized
oligonucleotides were analyzed by polyacrylamide gel
electrophoresis on denaturing gels and judged to be at least
85% full length material. The relative amounts of
phosphorothioate and phosphodiester linkages obtained in
synthesis were periodically checked by ³¹P nuclear magnetic
resonance spectroscopy, and for some studies oligonucleotides
were purified by HPLC, as described by Chiang et al., *J. Biol.*
Chem., 1991, 266, 18162-18171. Results obtained with HPLC-

- 49 -

purified material were similar to those obtained with non-HPLC purified material.

Example 7

Analysis of Oligonucleotide Inhibition of Novel Anti-apoptotic bcl-2-related Protein Expression

Antisense modulation of novel anti-apoptotic bcl-2-related protein expression can be assayed in a variety of ways known in the art. For example, mRNA levels can be quantitated by Northern blot analysis, RNase protection assay (RPA), competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, et al., *Current Protocols in Molecular Biology*, Volume 1, John Wiley & Sons, Inc., 1993, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, et al., *Current Protocols in Molecular Biology*, Volume 1, John Wiley & Sons, Inc., 1996, pp. 4.2.1-4.2.9. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions. Other methods of PCR are also known in the art. Probes and primers are designed to hybridize to the target nucleic acid sequence, using published sequence information (for example, Genbank accession no. U29680 for human A1 and Genbank accession no. L08246 for human mcl-1).

Novel anti-apoptotic bcl-2-related protein levels can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA, flow cytometry or fluorescence-activated cell sorting (FACS). Antibodies directed to bcl-2-related proteins can be identified and obtained from a variety of sources, such as PharMingen Inc., San Diego CA, or can be prepared via conventional antibody generation methods.

- 50 -

5 Methods for preparation of polyclonal antisera are taught in,
for example, Ausubel, et al., *Current Protocols in Molecular*
10 *Biology*, Volume 2, John Wiley & Sons, Inc., 1997, pp. 11.12.1-
11.12.9. Preparation of monoclonal antibodies is taught in,
5 for example, Ausubel, et al., *Current Protocols in Molecular*
Biology, Volume 2, John Wiley & Sons, Inc., 1997, pp. 11.4.1-
11.11.5.

15 Immunoprecipitation methods are standard in the art and
can be found at, for example, Ausubel, et al., *Current*
10 *Protocols in Molecular Biology*, Volume 2, John Wiley & Sons,
Inc., 1998, pp. 10.16.1-10.16.11. Western blot (immunoblot)
20 analysis is standard in the art and can be found at, for
example, Ausubel, et al., *Current Protocols in Molecular*
Biology, Volume 2, John Wiley & Sons, Inc., 1997, pp. 10.8.1-
25 10.8.21. Enzyme-linked immunosorbent assays (ELISA) are
standard in the art and can be found at, for example, Ausubel,
et al., *Current Protocols in Molecular Biology*, Volume 2, John
Wiley & Sons, Inc., 1991, pp. 11.2.1-11.2.22.

Example 8

20 RNase Protection Assay for Analysis of mRNA Levels

35 The ribonuclease (RNase) protection assay is a sensitive
and specific method for quantitating expression levels (Zinn,
et al., *Cell*, 1983, 34:865-79). The method is based on the
hybridization of a target RNA to an *in vitro* transcribed
40 25 ³²P-labeled anti-sense RNA probe from a DNA template. RNase
treatment follows, resulting in degradation of single-stranded
RNA and excess probe. The probe and target RNA are resolved
by denaturing polyacrylamide gel electrophoresis with the
45 "protected" probe visualized using autoradiography or beta
30 imaging equipment. Template sets can be purchased (PharMingen
Inc., San Diego, CA) which contain a series of biologically
relevant templates, each of distinct length and each
50 representing a sequence in a distinct mRNA species. Each

- 51 -

5 template set is capable of detecting up to 11 unique gene
messages in a single reaction mix in addition to one or more
housekeeping genes, L32 and GAPDH, which serve as internal
10 controls. These template sets allow for multiple
5 determinations to be made from a single sample. Multi-probe
RPA can be performed on total RNA preparations derived by
standard methods, without further purification of poly-A⁺ RNA.

15 Oligonucleotides were evaluated for their respective
effects on mcl-1 mRNA levels along with total bcl-x mRNA
10 levels, using the RIBOQUANT™ RNase protection kit (Pharmingen,
San Diego, CA). All assays were performed according to
20 manufacturer's protocols. Briefly, multi-probe DNA template
sets were used to generate antisense RNA transcripts
radiolabeled with dUTP-³²P. The template set used for
25 apoptosis genes was the human hAPO-2 set. These radiolabeled
probes were hybridized overnight with typically 10 µg of total
cellular RNA. The reaction mixture was then digested with
30 single-strand RNases to generate the protected fragments which
were electrophoresed through a 5% acrylamide/urea gel.
20 Protected bands were visualized and quantitated using a
PhosphorImager (Molecular Dynamics, Sunnyvale, CA).

35 Example 9

Antisense Inhibition of Human A1 Expression- Phosphorothioate Oligodeoxynucleotides

40 25 In accordance with the present invention, a series of
oligonucleotides were designed to target human A1 RNA, using
published sequences (Karsan et al., 1996, Blood 87, 3089-3096;
Genbank Accession no. U29680, incorporated herein as SEQ ID
45 NO: 1). The oligonucleotides are shown in Table 1.

50

55

Table 1
Antisense oligonucleotides targeted to human A1

"Nucl. pos." indicates the position of the first nucleotide on the target sequence (Genbank accession no. U29680; SEQ ID NO: 1) to which the oligonucleotide hybridizes. "PS" = 5' phosphorothioate linkage. "5meC" = 5-methylcytosine. "Deoxy" = 2'-H.

ISIS #	Sequence	Nucl. pos.	Target region	Chemistry	SEQ ID NO:	Activity (% of control)	% Inhib
17483	TGTGCTGAGAATGTCACATC	1	5' UTR	PS; deoxy; C = 5meC	2	91	09
17484	TTGAAGCTGTTGAGGCAATG	19	5' UTR	PS; deoxy; C = 5meC	3	55	45
17485	AAGTCTTGAGCTGGCTCACC	39	5' UTR	PS; deoxy; C = 5meC	4	51	49
17486	CTGTCACTCTCTGCCTGGTG	66	START CODON	PS; deoxy; C = 5meC	5	49	51
17487	CTGTAATATATCCAAATTC	91	CODING	PS; deoxy; C = 5meC	6	103	--
17488	GCAGATAGTCTGAGCCAGC	111	CODING	PS; deoxy; C = 5meC	7	52	48
17489	TGGACTGAGAACGCAACATT	193	CODING	PS; deoxy; C = 5meC	8	40	60
17490	AATAGTGTTCTGGCAGTGTG	271	CODING	PS; deoxy; C = 5meC	9	44	56
17491	TGATGCCGTCTTCAAACTCC	309	CODING	PS; deoxy; C = 5meC	10	45	55
17492	TGCTGTCGTAGAGTTTCTT	379	CODING	PS; deoxy; C = 5meC	11	48	52
17493	TCACAGATCTTCTGTTAAC	556	CODING	PS; deoxy; C = 5meC	12	44	56
17494	TGGAGTGTCTCTTCTGTGTC	604	STOP CODON	PS; deoxy; C = 5meC	13	22	78
17495	ATCGTTCCATATCATGTCAG	648	3' UTR	PS; deoxy; C = 5meC	14	24	76

5

10

15

20

25

30

35

40

45

50

55

17496	CAAATTTCCATAACTCTGG	724	3'UTR	PS; deoxy; C = 5meC	15	36	64
17497	CATACAATTTATTCAATTACA	750	3'UTR	PS; deoxy; C = 5meC	16	69	31

- 54 -

Oligonucleotides were tested by Northern blot analysis in human umbilical vein endothelial cells (HUVEC) at a concentration of 100 nM. HUVECs were grown to 80% confluency and washed three times with pre-warmed (37°C) Opto-MEM™ (Life Technologies, Inc., Gaithersburg, MD). Oligonucleotides were premixed with 10 µg/ml Lipofectin™ reagent (Life Technologies, Inc., Gaithersburg, MD) in Opti-MEM and applied to washed cells. Cells were incubated with oligonucleotide for 4 hr at 37°C after which the medium was removed and replaced with fresh medium. Total cellular RNA was isolated using an RNeasy™ kit (Qiagen Inc., Valencia, CA). Isolated RNA was separated on a 1% agarose/formaldehyde gel, transferred to a Hybond N+ nylon membrane (Amersham, Arlington Heights, IL) overnight, and UV-crosslinked in a Stratalinker 2400 (Stratagene, La Jolla, CA). Blots were hybridized for several hours with a single-stranded PCR ³²P-labeled probe generated with a primer having the sequence AGAAGTATGTGTTGGCAATCGT (SEQ ID NO: 17) according to published methods. Bednarczuk, T.A. et al., *BioTechniques*, 1991, 10, 478. PCR was used to amplify nucleotides 22 to 684 of the A1 sequence. Northern blots were stripped and reprobated with random-primed ³²P-labeled human G3PDH cDNA to confirm equivalent loading of RNA samples.

The results are shown in Table 1. As can be seen in the table, phosphorothioate oligodeoxynucleotides having SEQ ID NO: 4, 5, 7, 8, 9, 10, 11, 12, 13, 14 and 15 inhibited A1 mRNA expression by approximately 50% or more in this experiment. Of these, ISIS 17494 (SEQ ID NO: 13, targeting the stop codon) and ISIS 17495, SEQ ID NO: 14, targeting the 3' UTR just downstream of the stop codon, gave over 75% inhibition of A1 expression.

Example 10**Antisense Inhibition of A1 Expression- Mixed Backbone 2'-MOE
Gapmer Oligonucleotides**

A second series of oligonucleotides targeted to human A1 was synthesized. The oligonucleotides are shown in Table 2. Target sites are indicated by nucleotide numbers, as given in the sequence source reference to which the oligonucleotide binds.

All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-O-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) in the deoxy gap and phosphodiester (P=O) in the wings. Cytidine residues throughout the molecule are 5-methylcytidines. Oligonucleotides were tested by Northern blot analysis as described in the previous example. Results are shown in Table 2.

Table 2
Gapped chimeric oligonucleotides targeted to A1

"Nucl. pos." indicates the position of the first nucleotide on the target sequence (Genbank accession no. U29680; SEQ ID NO: 1) to which the oligonucleotide hybridizes. "N.D." indicates no data available. "PS" = phosphorothioate linkage; "PO" = phosphodiester linkage. "Deoxy" = 2'H; "2'MOE" = 2'-O-methoxyethyl; "5mec" = 5-methylcytosine.

ISIS #	Sequence	Nucl. pos.	Target region	Chemistry	SEQ ID NO:	Activity (% of control)	%inhb
17498	TGTGCTGAGATGCTCACTC	1	5' UTR	2'MOE/deoxy; PO/PS C = 5mec	2	64	36
17499	TTGAAGCTGTTGAGGCAATG	19	5' UTR	2'MOE/deoxy; PO/PS C = 5mec	3	25	75
17500	AAGTCTTGAGCTGGCTCACC	39	5' UTR	2'MOE/deoxy; PO/PS C = 5mec	4	44	56
17501	CTGTCAATCTTCTGCTGGTG	66	START CODON	2'MOE/deoxy; PO/PS C = 5mec	5	13	87
17502	CTGTAATATATCCAAATTC	91	CODING	2'MOE/deoxy; PO/PS C = 5mec	6	N.D.	
17503	GCAGATAGTCTGAGCCAGC	111	CODING	2'MOE/deoxy; PO/PS C = 5mec	7	24	76

17504	TGGACTGAGAACGCAACATT	193	CODING	2'MOE/deoxy; PO/PS C = 5meC	8	16	84
17505	AATAGTGTTCCTGGCAGTGTC	271	CODING	2'MOE/deoxy; PO/PS C = 5meC	9	11	89
17506	TGATGCCGTCTTCAAACTCC	309	CODING	2'MOE/deoxy; PO/PS C = 5meC	10	15	85
17507	TGCTGTCGTAGAAGTTTCTT	379	CODING	2'MOE/deoxy; PO/PS C = 5meC	11	9.3	90.7
17508	TCACAGATCTTTCCTGTAAC	556	CODING	2'MOE/deoxy; PO/PS C = 5meC	12	9.5	90.5
17509	TGGAGTGTCCTTCTGGTCA	604	STOP CODON	2'MOE/deoxy; PO/PS C = 5meC	13	6.7	93.3
17510	ATCGTTTCCATATCAGTCAG	648	3'UTR	2'MOE/deoxy; PO/PS C = 5meC	14	7.6	92.4
17511	CAAAATTCCATAACTCTGG	724	3'UTR	2'MOE/deoxy; PO/PS C = 5meC	15	15	85
17512	CATACAATTTATTCAATTACA	750	3'UTR	2'MOE/deoxy; PO/PS C = 5meC	16	22	78

- 58 -

As 2'-MOE gapmers, SEQ ID NO: 3, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15 and 16 gave over 75% reduction in A1 mRNA. Of these, SEQ ID NO: 11 (ISIS 17507), 12 (ISIS 17508), 13 (ISIS 17509) and 14 (ISIS 17510) gave over 90% reduction of A1 expression.

Dose-response experiments using ISIS 17510 (SEQ ID NO: 14, 2' MOE gapmer) gave an IC50 of less than 10 nM for A1 mRNA reduction using this compound.

Northern blot experiments using ISIS 17510 and a probe specific for the bcl-2 family member bcl-X demonstrated that the A1 antisense compound did not affect bcl-x mRNA levels.

Example 11

Antisense Inhibition of A1 Expression- Mixed Backbone 2'-MOE Hemimer Oligonucleotides

A third series of oligonucleotides targeted to human A1 was synthesized. The oligonucleotides are shown in Table 3. Target sites are indicated by nucleotide numbers, as given in the sequence source reference to which the oligonucleotide binds.

All compounds in Table 3 are chimeric "hemimer" oligonucleotides 20 nucleotides in length, composed of ten contiguous 2'-deoxynucleotides, joined to ten contiguous 2'-O-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) in the deoxy region and phosphodiester (P=O) in the 2'-MOE region. Cytidine residues throughout the molecule are 5-methylcytidines.

Oligonucleotides were tested by Northern blot analysis as described in previous examples. Results are shown in Table 3.

Table 3
 "Hemimer" chimeric oligonucleotides targeted to A1

"Nucl. pos." indicates the position of the first nucleotide on the target sequence (Genbank accession no. U29680; SEQ ID NO: 1) to which the oligonucleotide hybridizes. "PS" = 5' phosphorothioate linkage; "PO" = phosphodiester linkage. "Deoxy" = 2'H; "2'MOE" = 2'-O-methoxyethyl. "5meC" = 5-methylcytosine.

IS18 #	Sequence	Nucl. pos.	Target region	Chemistry	SEQ ID NO:	Activity (% of control)	% Inhib
17513	TGTGCTGAGAAATGCTCACTC	1	5' UTR	Deoxy/2'MOE,PS/PO C = 5meC	2	73	27
17514	TTGAAGCTGTTGAGGCAATG	19	5' UTR	Deoxy/2'MOE,PS/PO C = 5meC	3	40	60
17515	AAGTCTTGAGCTGGCTCACC	39	5' UTR	Deoxy/2'MOE,PS/PO C = 5meC	4	30	70
17516	CTGTCATCTTCTGCCCTGGTG	66	START CODON	Deoxy/2'MOE,PS/PO C = 5meC	5	32	68
17517	CTGTAATATATCCAAATTC	91	CODING	Deoxy/2'MOE,PS/PO C = 5meC	6	26	74
17518	GCAGATAGTCTCTGAGCCAGC	111	CODING	Deoxy/2'MOE,PS/PO C = 5meC	7	32	68
17519	TGGACTGAGAACGCAACATT	193	CODING	Deoxy/2'MOE,PS/PO C = 5meC	8	32	68

17520	AATAGTGTCTCTGGCAGTGTC	271	CODING	Deoxy/2'MOE, PS/PO C = 5meC	9	37	63
17521	TGATGCCGTCTTCAAACTCC	309	CODING	Deoxy/2'MOE, PS/PO C = 5meC	10	29	71
17522	TGCTGTCGTAGAAAGTTTCTT	379	CODING	Deoxy/2'MOE, PS/PO C = 5meC	11	23	77
17523	TCACAGATCTTCTCTGTAAC	556	CODING	Deoxy/2'MOE, PS/PO C = 5meC	12	16	84
17524	TGGAGTGTCCTTTCTGGTCA	604	STOP CODON	Deoxy/2'MOE, PS/PO C = 5meC	13	8.5	91.5
17525	ATCGTTTCCATATCAGTCAG	648	3' UTR	Deoxy/2'MOE, PS/PO C = 5meC	14	13	87
17526	CAAAATTTCATAAAGTCTGG	724	3' UTR	Deoxy/2'MOE, PS/PO C = 5meC	15	64	36
17527	CATACAATTATTCATTACA	750	3' UTR	Deoxy/2'MOE, PS/PO C = 5meC	16	47	53

- 61 -

As 2' MOE hemimers, SEQ ID NO: 11 (ISIS 17522), 12 (ISIS 17523), 13 (ISIS 17524) and 14 (ISIS 17525) gave at least 75% inhibition of A1 mRNA levels. Of these, ISIS 17524 (SEQ ID NO: 13) gave over 90% inhibition.

5 Example 12

15 Antisense Inhibition of Human mcl-1 Expression-Phosphorothioate 2'-moe Gapmer Oligonucleotides

In accordance with the present invention, a series of oligonucleotides were designed to target human mcl-1 RNA, using published sequences (Kozopas et al., Proc. Natl. Acad. Sci. USA., 1993, 90, 3516-3520, Genbank Accession No. L08246, incorporated herein as SEQ ID NO: 18). The oligonucleotides are shown in Table 4. All compounds in Table 4 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-O-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout. Cytidine residues throughout the molecule are 5-methylcytidines.

Oligonucleotides were tested by Northern blot analysis approximately as described in previous examples, using the human melanoma cell line C8161 (a gift of Dr. Dan Welch, Hershey Medical Center, Hershey, PA). Oligonucleotide (100 nM) and Lipofectin™ (10 µg/ml) were mixed and incubated for 30 minutes at room temperature. Cells were washed twice with Opti-MEM™, oligonucleotide mixture was added and cells were incubated at 37° for 4 hours. Oligonucleotide was replaced with fresh medium (RPMI with 10% FCS). 24 hours later, RNA was isolated using the Rneasy™ kit (Qiagen, Valencia, CA). RNA was electrophoresed, transferred to nitrocellulose and probed with a random-primed mcl-1 EST probe (ATCC 1064916

- 62 -

I.M.A.G.E. Clone ID: 796506), obtained from the American Type Culture Collection (Manassas, VA) and labeled with the Stripeasy™ kit (Ambion, Austin, TX).

Results are shown in Table 4.

Table 4
Antisense oligonucleotides targeted to human mcl-1
"Nucl. pos." indicates the position of the first nucleotide on the target sequence (Genbank accession no. U08246; SEQ ID NO:18) to which the oligonucleotide hybridizes. "N.D." indicates no data available. "PS" = phosphorothioate linkage; "PO" = phosphodiester linkage. "Deoxy" = 2'H; "2'MOE" = 2'-O-methoxyethyl. "5meC" = 5- methylcytosine.

ISIS #	Sequence	Nucl. pos.	Target region	Chemistry	SEQ ID NO.	Activity (% of control)	% Inhib
20404	GCCAAACATTGCCAGTCGCC	50		PS; 2'MOE/deoxy; C = 5 meC	19	83.2	16.8
20405	AGCCAAAAGTCGCCCTCCCG	152		PS; 2'MOE/deoxy; C = 5 meC	20	72.8	27.2
20406	CTCGTACCGTCCAGTCCT	431		PS; 2'MOE/deoxy; C = 5 meC	21	63.8	36.2
20407	TGTTATTACAGATTCCCG	501		PS; 2'MOE/deoxy; C = 5 meC	22	44.1	55.9
20408	TTGGCTTTGTCTCTTGCG	636		PS; 2'MOE/deoxy; C = 5 meC	23	38.4	61.6
20409	GAGAGTCACAATCCTGCCCC	842		PS; 2'MOE/deoxy; C = 5 meC	24	68.6	31.4
20410	AAAGCCAGCAGCACATTCT	1045		PS; 2'MOE/deoxy; C = 5 meC	25	66.6	33.4
20411	CCTCTTGCCACTTGCTTTTC	1230		PS; 2'MOE/deoxy; C = 5 meC	26	77.7	22.3

- 64 -

20412	CACAGGTCACTGGCATTCCTT	1519		PS; 2'MOE/deoxy; C = 5 meC	27	97.4	2.6
20413	AAGAATCATGGAACCAAGCC	1676		PS; 2'MOE/deoxy; C = 5 meC	28	84.1	15.9
20414	CTCTCAATCCAGGTTTCA	2091		PS; 2'MOE/deoxy; C = 5 meC	29	65.4	34.6
20415	GGTCAAAATGGAAGGAAGTCA	2231		PS; 2'MOE/deoxy; C = 5 meC	30	71.9	28.1
20416	CAATGTCTCTCCATCCACC	2366		PS; 2'MOE/deoxy; C = 5 meC	31	37.5	62.5
20417	AAATCCAAAGATGCCAATGC	2564		PS; 2'MOE/deoxy; C = 5 meC	32	159.5	--
20418	CAGTGCCAAATCTAAAGG	2722		PS; 2'MOE/deoxy; C = 5 meC	33	87.5	12.5
20419	CTTCCTCCACCTCTCAATG	2866		PS; 2'MOE/deoxy; C = 5 meC	34	N.D.	
20420	GGCAGTTCTTCCCCATTACA	3207		PS; 2'MOE/deoxy; C = 5 meC	35	43.8	56.2
20421	ATTTGGCAGACAGGCTTTTA	3391		PS; 2'MOE/deoxy; C = 5 meC	36	50.6	49.4
20422	TAGACCACCTGCCTCCTCCT	3591		PS; 2'MOE/deoxy; C = 5 meC	37	62.5	37.5
20423	GTCCTAACCCCTTCTTGGCAG	3821		PS; 2'MOE/deoxy; C = 5 meC	38	68.8	31.2

- 65 -

SEQ ID NO: 22 (ISIS 20407), 23 (ISIS 20408), 31 (ISIS 20416), 35 (ISIS 20420), and 36 (ISIS 20421) inhibited mcl-1 mRNA levels by approximately 50% or more. Of these, ISIS 20408 and 20416 showed greatest reduction of mcl-1 expression.

A dose-response experiment was done using ISIS 20407, 20408 and 20416. RNase protection assay was used to detect mcl-1 mRNA levels, using the hApo2 probe set (Pharmingen, San Diego, CA). ISIS 20407 demonstrated an IC₅₀ of approximately 100 nM. ISIS 20408 and 20416 demonstrated IC₅₀s of approximately 25 nM.

Example 13

Scid-human Leukemia Xenograft Model and Measurement of Apoptosis in Xenografts

10⁷ SEM-K2 cells in exponential phase of growth are injected subcutaneously into 8 SCID-NOD mice as a bolus (suspended in sterile saline). Engraftment and tumor formation occurs over a 2-3 week period. Micro Alzet pumps (Alza, Newark, DE) capable of delivering a continuous subcutaneous infusion over 14 days are used to deliver a dose of 100 µg per day (equivalent to 5 mg/kg) of antisense oligonucleotide into three animals. The remaining two animals received vehicle (sterile saline) only.

The expression of target protein measured in SEM-K2 cells from SCID-hu xenografts is measured using quantitative flow cytometry.

Xenografts are removed after sacrifice and mechanically dispersed into large volumes of medium. Leukocytes are purified by density gradient centrifugation and washed with medium before resuspending in 1 ml volumes at 1-5 x 10⁶ cells/ml. Cells are incubated at 37° C in 95% humidified air/5% CO₂ for 2 hours prior to induction of apoptosis with 20 µg/ml VP16 (Etoposide) over 24 hours. Apoptosis is assessed nonspecifically using quantification of light scatter changes;

- 66 -

reduction in side scatter (due to chromatin condensation) and reduction in forward scatter (due to cell shrinkage) are early changes associated with apoptosis. Bimodal population distributions consisting of apoptotic and non-apoptotic cells can be measured respectively allowing estimation of an apoptotic index for treated and negative control. Fold increase in apoptosis is calculated from their ratio. More specific determination of apoptosis is achieved using the Apo-Alert Caspase-3 Colorimetric Assay Kit (Clontech, Palo Alto, CA). This is a DEVD-specific caspase assay, a quantitative assay for the activity of caspase-3, a member of the caspase family thought to mediate apoptosis in most mammalian cell types. This assay utilizes a synthetic tetrapeptide, Asp-Glu-Val-Asp (DEVD) (SEQ ID NO: 39), labeled with either a fluorescent molecule, 7-amino-4-trifluoromethyl coumarin (AFC), or a colorimetric molecule, p-nitroanilide (pNA) as substrates. DEVD-dependent protease activity is assessed by detection of the free AFC or pNA cleaved from the substrates. Cell lysates are incubated with DEVD conjugated to paranitroanilide, a colorimetric substrate cleaved by CPP32 (caspase-3) and detectable using colorimetric spectrophotometry at 405 nm. The fold increase in OD_{405nm} is used to determine the net VP16-induced apoptosis.

Example 14

Western blot analysis of mcl-1 protein levels in C8161 cells

Western blot analysis (immunoblot analysis) was carried out using standard methods. C8161 cells were transfected in 5 mL volumes at approximately 8×10^5 cells/10 cm plate. 200 nM of ISIS 20408 (SEQ ID NO: 23) or 8-basepair mismatch control ISIS 105232; TGGGTCTGGTTTCTCTGTCG; SEQ ID NO: 40) was added to cells with cationic lipid. Cells were analyzed at fixed intervals after treatment.

For Western blot analysis, cells were washed once in phosphate-buffered saline and pelleted by centrifugation at

- 67 -

5 1200 rpm for 5 minutes, and resuspended in cell lysis buffer
(5M NaCl, 0.1 M HEPES, 500 mM sucrose, 0.5M EDTA, 100 mM
spermine, 1 mg/ml aprotinin, 10% Triton X-100) for 15 minutes
10 on ice. Total protein was quantified spectrophotometrically
(BioRad) and 100 µg lysate was loaded onto 15% polyacrylamide
5 gels and run at 200V for 45 minutes. Following protein
transfer to nitrocellulose membrane, blots were immunostained
15 with primary mcl-1 antibody (Santa Cruz Biotechnology, Inc.,
Santa Cruz CA) followed by horseradish peroxidase-conjugated
20 goat anti-mouse secondary antibody (Santa Cruz Biotechnology,
Inc., Santa Cruz CA). Protein was qualitatively visualized by
ECL (Amersham, Piscataway NJ) and exposure on photographic
film.

ISIS 20408 was tested for the ability to reduce mcl-1
25 protein levels in C8161 cells, and was found to reduce mcl-1
protein levels by 90% after 4 hours, at an oligonucleotide
dose of 200nM.

30 Example 15

Ultraviolet Irradiation of C8161 cells

20 C8161 cells were irradiated with UV-B light when they
were approximately 70-80% confluent or 24 hours after
35 treatment with oligonucleotide. Immediately before UV-B
treatment, cells were washed twice with PBS and then exposed
to UV-B light in a Stratalinker UV Crosslinker 1800 model
25 (Stratagene, La Jolla CA) containing 5 15-watt 312 nm bulbs.
The cells were exposed to 0 mJ/m², 5 mJ/m² or 10 mJ/m² of UV-B
40 radiation. The dose of UV-B radiation was calibrated using a
UVX radiometer (UVP). Following UV irradiation, the cells were
incubated in the standard medium for an additional 24 hours.
45 Control plates for UV-B treatment were simply washed 3 times
in PBS and incubated in the standard medium for an additional
24 hours. Cells were examined for apoptosis by staining the
50 ethanol-fixed cell nuclei with propidium iodide and examining
the DNA content by flow cytometry. Apoptotic cells were

- 68 -

identified by their sub-diploid DNA content. Cells were washed twice with cold PBS and resuspended in 1 ml of 70% ethanol. After 1 hour incubation at room temperature, cells were washed in PBS and resuspended in 1 ml propidium iodide staining solution (50 μ g/ml propidium iodide, 0.5 U/ml RNase A, 2000 U/ml RNase T1. After 30 minutes at room temperature, cell cycle analysis was performed by flow cytometry using a Becton Dickinson Calibur FACS analyzer. The fluorescence of individual nuclei of 10,000 cells was measured using a FACScan flow cytometer. Results were expressed as percentage apoptotic cells.

Example 16**Antisense sensitization of C8161 cells to UV-induced cell death**

C8161 cells were treated with 200 nM ISIS 20408 or the 8-basepair mismatch control (ISIS 105232; SEQ ID NO: 40) and exposed to ultraviolet (UV) radiation as described in the previous example. The percent apoptotic cells was quantitated by propidium iodide staining according to standard methods. Results are shown in Table 5.

Table 5**Combination of ISIS 20408 and UV irradiation**

Compound	UV mJ/M ²	% Apoptotic cells (approx)	SEQ ID NO:
No oligo	0	0.2	--
	5	1.0	--
	10	5.2	--
ISIS 20408	0	2	23
	5	4.7	" "
	10	11.5	" "
ISIS 105232	0	0.5	40
	5	1.5	" "

- 69 -

	10	6.3	" "
--	----	-----	-----

Thus the antisense treatment resulted in increased apoptosis;
i.e., sensitization to UV-induced apoptosis.

Example 17

Cisplatinium Treatment of C8161 Cells

Cisplatinium is an alkylating agent that causes DNA damage and can induce apoptosis. Gill and Windebank, 1998, J. Clin. Invest. 101:2842-2850. C8161 cells were treated with cisplatinium when they were approximately 70-80% confluent or 24 hours after treatment with oligonucleotide. Cis-diamminedichloroplatinum II (Cisplatinium, Sigma, St. Louis MO) was dissolved in distilled water at a concentration of 1 mg/ml, and was added to the standard medium at a dose range of 2.5 to 10 μ g/ml and incubated with cells for 24 hours.

Example 18

Antisense sensitization of C8161 cells to cisplatinium-induced cell death

C8161 cells were treated with 200 nM ISIS 20408 or the 8-basepair mismatch control ISIS 105232 and cisplatinium at various doses. The percent apoptotic cells was quantitated by propidium iodide staining according to standard methods as described in previous examples. Results are shown in Table 6.

Table 6

Combination of ISIS 20408 and Cisplatinium

Compound	Cisplatinium dose (μ g/ml)	% Apoptotic cells (approx)	SEQ ID NO:
No oligo	0	2	--
	2.5	5	--
	5	11	--
	10	30	--

- 70 -

ISIS 20408	0	11.5	23
	2.5	39	" "
	5	40	" "
	10	80	" "
ISIS 105232	0	4	40
	2.5	11.5	" "
	5	17.5	" "
	10	52	" "

Thus the cells have been sensitized to the apoptotic stimulus
(in this case the cytotoxic chemotherapeutic drug cisplatinium)
5 after antisense treatment resulting in increased apoptosis.

Claims

5

10

THIS PAGE BLANK (USPTO)

15

20

25

30

35

40

45

50

55

- 71 -

What is claimed is:

1. An antisense compound 8 to 30 nucleotides in length targeted to a nucleic acid molecule encoding a novel anti-apoptotic bcl-2-related protein, wherein said antisense compound modulates the expression of said novel anti-apoptotic bcl-2-related protein.
2. The antisense compound of claim 1 which is an antisense oligonucleotide.
3. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.
4. The antisense compound of claim 3 wherein the modified internucleoside linkage of the antisense oligonucleotide is a phosphorothioate linkage.
5. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.
6. The antisense compound of claim 5 wherein the modified sugar moiety of the antisense oligonucleotide is a 2'-O-methoxyethyl sugar moiety.
7. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.
8. The antisense compound of claim 7 wherein the modified nucleobase is a 2'-O-methoxyethyl modified cytosine.

- 72 -

5
10
9. The antisense compound of claim 8 wherein the modified nucleobase of the antisense oligonucleotide is a 5-methylcytosine.

15
10. The antisense compound of claim 1 which is a 5 chimeric oligonucleotide.

20
11. The antisense compound of claim 1 which is targeted to a nucleic acid encoding human A1.

25
12. The antisense compound of claim 11 wherein the antisense oligonucleotide comprises SEQ ID NO: 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16.

30
13. The antisense compound of claim 1 which is targeted to a nucleic acid encoding human mcl-1.

35
14. The antisense compound of claim 13 wherein the antisense oligonucleotide comprises SEQ ID NO: 22, 23, 31, 35, 15 or 36.

40
15. A pharmaceutical composition comprising the antisense compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

45
16. The pharmaceutical composition of claim 15 further comprising a colloidal dispersion system.

50
17. The pharmaceutical composition of claim 15 further comprising a chemotherapeutic agent for the treatment of cancer.

55
18. A method of inhibiting the expression of a novel anti-apoptotic bcl-2-related protein in cells or tissues comprising contacting said cells or tissues with the antisense

- 73 -

compound of claim 1 so that expression of said novel anti-apoptotic bcl-2-related protein is inhibited.

19. The method of claim 18 wherein the antisense compound is targeted to a nucleic acid molecule encoding human A1.

20. The method of claim 18 wherein the antisense compound is targeted to a nucleic acid molecule encoding human mcl-1.

21. A method of promoting apoptosis in human cells or tissues comprising contacting said cells or tissues with the antisense compound of claim 1 so that expression of said novel anti-apoptotic bcl-2-related protein is inhibited.

22. The method of claim 21 wherein the antisense compound is targeted to a nucleic acid molecule encoding human A1.

23. The method of claim 21 wherein the antisense compound is targeted to a nucleic acid molecule encoding human mcl-1.

24. A method of treating a human having a disease or condition associated with a novel anti-apoptotic bcl-2-related protein comprising administering to said animal a therapeutically or prophylactically effective amount of the antisense compound of claim 1 so that expression of said novel anti-apoptotic bcl-2-related protein is inhibited.

25. The method of claim 24 wherein the antisense compound is targeted to a nucleic acid molecule encoding human A1.

5

- 74 -

10

26. The method of claim 24 wherein the antisense compound is targeted to a nucleic acid molecule encoding human mcl-1.

15

27. A method of treating a human having a disease or condition characterized by a reduction in apoptosis comprising administering to said human a prophylactically or therapeutically effective amount of the antisense compound of claim 1.

20

28. The method of claim 27 wherein the antisense compound is targeted to a nucleic acid molecule encoding human A1.

25

29. The method of claim 27 wherein the antisense compound is targeted to a nucleic acid molecule encoding human mcl-1.

30

30. A method of treating a human having a disease or condition for which promotion of apoptosis is desired comprising administering to said human a prophylactically or therapeutically effective amount of the antisense compound of claim 1.

35

40

31. The method of claim 30 wherein the antisense compound is targeted to a nucleic acid molecule encoding human A1.

45

32. The method of claim 30 wherein the antisense compound is targeted to a nucleic acid molecule encoding human mcl-1.

25

50

33. A method of treating cancer in a patient comprising:

55

- 75 -

5 (a) administering to the patient a pharmaceutical composition of claim 15; and

10 (b) administering to the patient a chemotherapeutic agent for the treatment of cancer.

5 34. The pharmaceutical composition of claim 17 wherein the chemotherapeutic agent for the treatment of cancer is cisplatin.

20 35. A method of sensitizing human cells or tissues to apoptosis comprising contacting said cells or tissues with the antisense compound of claim 1 so that expression of said novel anti-apoptotic bcl-2-related protein is inhibited.

25 36. The method of claim 35 wherein said antisense compound is administered in combination with an apoptotic stimulus.

30 15 37. The method of claim 36 wherein the apoptotic stimulus is radiation or a chemotherapeutic agent.

35 38. The method of claim 37 wherein the chemotherapeutic agent is cisplatin.

40 39. A method of sensitizing human cells or tissues to an apoptotic stimulus comprising contacting said cells or tissues with the antisense compound of claim 1 so that expression of said novel anti-apoptotic bcl-2-related protein is inhibited.

45 40. The method of claim 39 wherein the apoptotic stimulus is radiation or a chemotherapeutic agent.

50 41. The method of claim 40 wherein the chemotherapeutic agent is cisplatin.

55

SEQUENCE LISTING

<110> Ackermann, Elizabeth J.
 Bennett, C. Frank
 Dean, Nicholas M.
 Marcusson, Eric G.
 Isis Pharmaceuticals, Inc.

<120> Antisense Modulation of Novel Anti-apoptotic bcl-2-Related
 Proteins

<130> ISPH-0432

<140>
 <141>

<150> 09/226,568
 <151> 1999-01-07

<160> 40

<170> Patent In Ver. 2.0

<210> 1
 <211> 780
 <212> DNA
 <213> Homo sapiens

<400> 1

gagtgagcat	tctcagcaca	ttgcctcaac	agcttcaagg	tgagccagct	caagactttg	60
ctctccacca	ggcagaagat	gacagactgt	gaatttggat	atatttacag	gctggctcag	120
gactatctgc	agtgcgctct	acagatacca	caacctggat	caggtccaag	caaaacgtcc	180
agagtgtctac	aaaatgtttgc	gttctcagtc	caaaaagaag	tggaaaagaa	tctgaagtca	240
tgcttggaca	atgttaatgt	tgtgtccgta	gacactgcca	gaacactatt	caaccaagtg	300
atggaaaagg	agtttgaaga	cggcatcatt	aactggggaa	gaattgtaac	catatttgca	360
tttgaaggta	ttctcatcaa	gaaacttcta	cgacagcaaa	ttgccccgga	tgtggatacc	420
tataaggaga	tttcatattt	tgttgccggag	ttcataatga	ataacacagg	agaatggata	480
aggcaaaacg	gaggctggga	aaatggcctt	gtaaagaagt	ttgaacctaa	atctggctgg	540
atgacttttc	tagaagttac	aggaaagatc	tgtgaaatgc	tatctctcct	gaagcaatac	600
tgttgaccag	aaaggacact	ccatattgtg	aaaccggcct	aatttttctg	actgatatgg	660
aaacgattgc	caacacatac	ttctactttt	aaataaacia	ctttgatgat	gtaacttgac	720
cttccagagt	tatggaattt	ttgtcccat	gtaatgaata	aattgtatgt	atttttctct	780

<210> 2
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: antisense sequence

<400> 2
 tgtgctgaga atgctcactc 20

<210> 3
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: antisense sequence

<400> 3
ttgaagctgt tgaggcaatg 20

<210> 4
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 4
aagtccttgag ctggctcacc 20

<210> 5
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 5
ctgtcatctt ctgcctggtg 20

<210> 6
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 6
ctgtaaatat atccaaattc 20

<210> 7
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 7
gcagatagtc ctgagccagc 20

<210> 8
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 8
tggactgaga acgcaacatt 20

<210> 9
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 9
aatagtgttc tggcagtgtc 20

<210> 10
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 10
tgatgccgtc ttcaaactcc 20

<210> 11
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 11
tgctgtcgta gaagtttctt 20

<210> 12
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 12
tcacagatct ttctgtaac 20

<210> 13
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 13
tggagtgtcc tttctggcca 20

<210> 14
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 14
atcgtttcca tatcagtcag 20

<210> 15

<211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: antisense sequence

 <400> 15
 caaaatttcc ataactctgg 20

 <210> 16
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: antisense sequence

 <400> 16
 catacaattt attcattaca 20

 <210> 17
 <211> 22
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: antisense sequence

 <400> 17
 agaagtatgt gttggcaatc gt 22

 <210> 18
 <211> 3934
 <212> DNA
 <213> Homo sapiens

 <220>
 <223> Description of Artificial Sequence: antisense sequence

 <400> 18
 tccagtaagg agtcgggggtc ttccccagtt ttctcagcca ggccggcgccg gcgactggca 60
 atgtttggcc tcaaaagaaa cgcggtaatc ggactcaacc tctactgtgg gggggccggc 120
 ttgggggccc gcagcggccg cgcacccgc cgggagggc gacttttggc tacggagaag 180
 gaggcctcgg cccggcgaga gataggggga ggggagggc gcgcgggtgat tggcggaagc 240
 gccggcgcaa gcccccgcgc caccctcacg ccagactccc ggagggtcgc gcggccgccg 300
 cccattggcg ccgaggtccc cgacgtcacg gcgacccccg cgaggctgct tttcttcgcg 360
 cccacccgcc gcgcggcgcc gcttgaggag atggaagccc cggccgctga cgccatcatg 420
 tcgcccgaag aggagctgga cgggtacgag ccggagcctc tcgggaagcg gccggctgtc 480
 ctgcccgtgc tggagtgtgt cggggaatct ggtaataaca ccagtacgga cgggtcacta 540
 ccctcgacgc cgcgcgcagc agaggaggag gaggacgagt tgtaccggca gtcgctggag 600
 attatctctc ggtaccttcg ggagcaggcc accggcgcca aggcacacaa gccaatgggc 660
 aggtctgggg ccaccagcag gaaggcgtg gagaccttac gacgggttgg ggatggcgtg 720
 cagcgcaacc acgagacggt cttccaaggc atgcttcgga aactggacat caaaaacgaa 780
 gacgatgtga aatcggtgtc tcgagtgtat atccatgttt tcagcgacgg cgtaacaaac 840
 tggggcagga ttgtgactct catttctttt ggtgcctttg tggctaaaca cttgaagacc 900
 ataaaccaag aaagctgcat cgaaccatta gcagaaagta tcacagacgt tctcgtagg 960
 acaaacggg actggctagt taaacaaaga ggctgggatg ggtttgtgga gttcttccat 1020
 gtagaggacc tagaaggtgg catcaggaat gtgctgctgg cttttgcagg tgttgcctga 1080
 gtaggagctg gtttggcata tctaataaga tagccttact gtaagtgcga tagttgactt 1140
 ttaaccaacc accaccacca ccaaaaccag tttatgcagt tggactccaa gctgtaactt 1200
 cctagagttg caccctagca acctagccag aaaagcaagt ggcaagagga ttatggctaa 1260
 caagaataaa tacatgggaa gagtgtctcc cattgattga agagtcactg tctgaaagaa 1320

```

gcaagttca gtttcagcaa caaacaaact ttgtttggga agctatggag gaggactttt 1380
agatttagtg aagatggtag ggtggaaaga cttaatttcc ttgttgagaa caggaaagtg 1440
gccagtagcc aggcaagtc tagaattgat taccgcccga attcattaat ttactgtagt 1500
agtgttaaga gaagcactaa gaatgccagt gacctgtgta aaagttacaa gtaatagaac 1560
tatgactgta agcctcagta ctgtacaagg gaagcttttc ctctctctaa ttactgttcc 1620
cagtatactt cttagaaagt ccaagtgttc aggactttta taccgtgtat actttggctt 1680
ggttccatga ttcttacttt attagcctag ttatcacca ataactctg acggaaggct 1740
cagtaattag ttatgaatat ggatccctc aattcttaag acagcttgta aatgtatttg 1800
taaaaattgt atatatTTTT acagaaagtc tatttccttg aaacgaagga agtatcgaat 1860
ttacattagt ttttttcata cccttttgaa ctttgcaact tccgtaatta ggaacctgtt 1920
tcttacagct tttctatgct aaactttgtt ctgttcagtt ctagagtgtat tacagaacga 1980
attgatgtgt aactgtatgc agactgggtg tagtggaaca aatctgataa ctatgcaggt 2040
ttaaattttc ttatctgatt ttggttaagta ttccttagat aggttttctt tgaaaacctg 2100
ggattgagag gttgatgaat ggaaattctt tcacttcatt atatgcaagt tttcaataat 2160
taggtctaa gtaggtttta aggttactga tgacttaca ataatgggct ctgattgggc 2220
aatactcatt tgagttcctt ccatttgacc taatttaact ggtgaaattt aaagtgaatt 2280
catgggctca tctttaagc ttttactaaa agattttcag ctgaatggaa ctcatagct 2340
gtgtgcata aaaaagatca catcaggtgg atggagagac atttgatccc ttgtttgctt 2400
aataaattat aaaaatgatg cttggaaaag caggctagtc taacctgggt gctattatta 2460
ggcttgcttg ttacacacac aggtcctaag ctagtatgtc aataaagcaa atacttactg 2520
ttttgtttct attaatgatt cccaaacctt gttgcaagtt ttgtcattgg catctttgga 2580
tttcagctct gatgtttgtt ctatcagact taacctttta ttctctgtcc ttcttgaaa 2640
ttgtgatgg ttctgtctcc tctacagata ttatatcaa ttcctacagc tttccctgc 2700
catccctgaa ctctttctag ccctttttaga ttttggcact gtgaaacccc tgctggaaac 2760
ctgagtgacc ctccctcccc accaagagtc cacagacctt tcatctttca cgaacttgat 2820
cctgttagca ggtggtaata ccatgggtgc tgtgacacta acagtcattg agagggtgga 2880
ggaaagtcct tttccttgga ctgggtatctt tcaactatt gttttatcct gtctttgggg 2940
gcaatgtgtc aaaagtcctc tcaggaattt tcagaggaaa gaacatttta tgaggctttc 3000
tctaaagttt cttttgtata ggagtatgct cacttaaat tacagaaaga ggtgagctgt 3060
gttaaacctc agagtttaaa agctactgat aaactgaaga aagtgtctat attggaacta 3120
gggtcatttg aaagcttcag tctcggaaca tgaccttag tctgtggact ccatttaaaa 3180
ataggtatga ataagatgac taagaatgta atggggaaga actgcccctgc ctgcccctct 3240
cagagccata aggtcatctt tgctagagct atttttacct atgtatttat cgttcttgat 3300
cataagccgc ttatttatat catgtatctc taaggaccta aaagcacttt atgtagtttt 3360
taattaatct taagatctgg ttacggtaac taaaagcctg tctgccaaat ccagtggaaa 3420
caagtgcata gatgtgaatt ggtttttagg ggccccactt cccaattcat taggtatgac 3480
tgtggaata cagacaagga cttagttgat attttgggct tggggcagtg agggcttagg 3540
acaccccaag tggtttggga aaggagggag gagtgggtgg tttatagggg agggaggggc 3600
aggtgtgtcta agtgctgact ggctacgtag ttcgggcaaa tccctccaaa gggaaaggga 3660
ggatttgctt agaaggatgg ggctcccagt gactactttt tgacttctgt ttgtcttacg 3720
cttctctcag ggaaaaacat gcagtcctct agtgtttcat gtacattctg tggggggtgac 3780
acaccttggt tctgggttaa cagctgtact ttgtatagct gtgccaggaa gggttaggac 3840
caactacaaa ttaatgttgg ttgtcaaatg tagtgtgttt ccctaacttt ctgtttttcc 3900
tgagaaaaaa aaataaatct tttattcaaa taaa 3934

```

<210> 19
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 19
gccaaacatt gccagtcgcc

20

<210> 20
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: antisense sequence

<400> 20
agccaaaagt cgcctccccg 20

<210> 21
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 21
ctcgtaccog tccagtcct 20

<210> 22
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 22
tgttattacc agattccccg 20

<210> 23
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 23
ttggctttgt gtccttggcg 20

<210> 24
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 24
gagagtcaca atcctgcccc 20

<210> 25
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 25
aaagccagca gcacattcct 20

<210> 26
<211> 20
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: antisense sequence

<400> 26

cctccttgcca ctgcttttc

20

<210> 27

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: antisense sequence

<400> 27

cacaggtcac tggcattctt

20

<210> 28

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: antisense sequence

<400> 28

aagaatcatg gaaccaagcc

20

<210> 29

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: antisense sequence

<400> 29

ctctcaatcc caggttttca

20

<210> 30

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: antisense sequence

<400> 30

ggtcaaattg aaggaactca

20

<210> 31

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: antisense sequence

<400> 31

caaatgtctc tccatccacc

20

<210> 32
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 32
aaatccaaag atgccaatgc 20

<210> 33
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 33
cagtgccaaa atctaaaagg 20

<210> 34
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 34
cttcctccca cctctcaatg 20

<210> 35
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 35
ggcagttctt cccattaca 20

<210> 36
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 36
atttggcaga caggctttta 20

<210> 37
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 37
tagaccacct gcctcctcct 20

<210> 38
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 38
gtcctaacc ttcctggcac 20

<210> 39
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic peptide

<400> 39
Asp Glu Val Asp
1

<210> 40
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: antisense sequence

<400> 40
tgggtctggt ttctctgtcg 20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/29593

A. CLASSIFICATION OF SUBJECT MATTERIPC(7) : C07H 21/04, 21/02; C12Q 1/68, 15/63; A61K 48/00
US CL : 536/23.1, 24.3, 24.5; 435/6, 91.1, 375, 440; 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHEDMinimum documentation searched (classification system followed by classification symbols)
U.S. : 536/23.1, 24.3, 24.5; 435/6, 91.1, 375, 440; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,583,034 A (GREEN et al) 10 December 1996 (10.12.1996), column 2, line 50-67.	1-11, 13, 21-23
Y	CHAO J. R. mcl-1 Is an Immediate-Early Gene Activated by the Granulocyte-Macrophage Colony-Stimulating Factor(GM-CSF) Signaling Pathway and Is One Component of the GM-CSF Viability Response. Molecular and Cellular Biology. August 1998, Vol. 18, No. 8, pages 4883-4898, especially page 4894.	1-11, 13, 21-23
A	CROOKE S. T. Basic Principles of Antisense Therapeutics. In: Antisense Research And Applications, Chapter 1, Springer-Verlag Press, Berlin, Heidelberg, New York July 1998 pages 1-50, especially page 2-3	1-41

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" documents referring to an oral disclosure, use, exhibition or other means	"A" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

11 February 2000 (11.02.2000)

Date of mailing of the international search report

29 FEB 2000

Name and mailing address of the ISA/US

Comptroller of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

George Elliott

Telephone No. 703-308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/29593

Continuation of B. **FIELDS SEARCHED** Item 3: USPAT, EPO, JPO, CaPlus
search terms: antisense, ribozyme, aptamer, triplex, bcl7, bcl-2, apoptosis